



Contents lists available at ScienceDirect

## Journal of Number Theory

[www.elsevier.com/locate/jnt](http://www.elsevier.com/locate/jnt)

# On multiple $\mathbb{Z}_p$ -extensions of imaginary abelian quartic fields

Tsuyoshi Itoh<sup>1</sup>

College of Science and Engineering, Ritsumeikan University, 1-1-1 Noji Higashi, Kusatsu, Shiga 525-8577, Japan

## ARTICLE INFO

## Article history:

Received 26 February 2005

Revised 10 May 2010

Accepted 1 August 2010

Available online 6 September 2010

Communicated by David Goss

## MSC:

11R23

## Keywords:

Iwasawa modules

Multiple  $\mathbb{Z}_p$ -extensions

## ABSTRACT

Let  $k$  be an imaginary abelian quartic field and  $p$  an odd prime which splits completely in  $k$ . We give a sufficient condition for the validity of Greenberg's Generalized Conjecture (for multiple  $\mathbb{Z}_p$ -extensions) for  $k$  and  $p$ .

© 2010 Elsevier Inc. All rights reserved.

## 1. Introduction

First, we shall briefly explain about Iwasawa theory for multiple  $\mathbb{Z}_p$ -extensions (for details, see [5]). Let  $k$  be an algebraic number field and  $p$  a prime number. Let  $K/k$  be a  $\mathbb{Z}_p^d$ -extension (where  $d$  is a positive integer). We denote by  $L(K)$  the maximal unramified abelian pro- $p$  extension of  $K$ . Then it is known that the Galois group  $\text{Gal}(L(K)/K)$  is a finitely generated torsion module over the complete group ring  $\mathbb{Z}_p[[\text{Gal}(K/k)]]$ . Since  $\mathbb{Z}_p[[\text{Gal}(K/k)]]$  is isomorphic to the  $d$ -variable power series ring  $\Lambda_d := \mathbb{Z}_p[[T_1, T_2, \dots, T_d]]$ , we may regard an element of  $\mathbb{Z}_p[[\text{Gal}(K/k)]]$  as in  $\Lambda_d$  (under a fixed isomorphism).

A finitely generated  $\mathbb{Z}_p[[\text{Gal}(K/k)]]$ -module  $M$  is called pseudo-null if there are nontrivial two annihilators  $f, g \in \mathbb{Z}_p[[\text{Gal}(K/k)]]$  of  $M$  such that  $f$  and  $g$  are relatively prime (cf. [8]). For a finitely generated torsion  $\mathbb{Z}_p[[\text{Gal}(K/k)]]$ -module  $M$ , it is known that  $M$  is pseudo-null if and only if

E-mail address: [tsuyoshi.itoh@it-chiba.ac.jp](mailto:tsuyoshi.itoh@it-chiba.ac.jp).

<sup>1</sup> Current address: Division of Mathematics, Education Center, Faculty of Social Systems Science, Chiba Institute of Technology, 2-1-1 Shibazono, Narashino, Chiba 275-0023, Japan.

$\text{Ext}_{\mathbb{Z}_p[[\text{Gal}(K/k)]]}^1(M, \mathbb{Z}_p[[\text{Gal}(K/k)]])$  is trivial. (This fact follows from, e.g., [20, p. 16, Proposition 8].) When  $d = 1$ ,  $M$  is pseudo-null if and only if the order of  $M$  is finite (see, e.g., [24, Lemma 15.17]). Let  $\tilde{k}$  be the composite of all  $\mathbb{Z}_p$ -extensions of  $k$ . It is known that  $\tilde{k}/k$  is a  $\mathbb{Z}_p^d$ -extension with some positive integer  $d$  (see, e.g., [24, Theorem 13.4]). In this case, the following conjecture is known. It is often called Greenberg's Generalized Conjecture (GGC).

**Conjecture.** (See Greenberg [8].) *The Galois group  $\text{Gal}(L(\tilde{k})/\tilde{k})$  is pseudo-null as a module over  $\mathbb{Z}_p[[\text{Gal}(\tilde{k}/k)]]$  for every algebraic number field  $k$  and every prime number  $p$ .*

If  $k$  is totally real and Leopoldt's conjecture holds for  $k$  and  $p$ , GGC is equivalent to (usual) Greenberg's Conjecture (see [6,8]). We shall state some of known results for the case that  $k$  is not totally real. When  $k$  is an imaginary quadratic field, Minardi [16,17] showed that if  $p$  does not divide the class number of  $k$  then GGC holds for  $k$  and  $p$ . The author [10] gave some families of imaginary quadratic fields which satisfy GGC for  $p = 2$ . When  $k$  is an imaginary biquadratic field and  $p$  is an odd prime number which splits at most two distinct primes in  $k$ , Bandini [1] gave some sufficient conditions for which GGC holds. See also [2]. When  $k$  is the  $p$ th cyclotomic field, it is known that GGC holds for  $k$  and  $p$  if  $p < 1000$  (see McCallum and Sharifi [15], Sharifi [21,22]).

In the present paper, we shall give a sufficient condition for the case that the base field  $k$  is an imaginary abelian quartic field (namely, an imaginary cyclic quartic field or an imaginary biquadratic field) and  $p$  is an odd prime number which splits completely in  $k$ . The following is our main theorem.

**Theorem 1.1.** *Let  $k$  be an imaginary abelian quartic field and  $p$  an odd prime number which splits completely in  $k$ . Assume that the class number of  $k$  is not divisible by  $p$ . We denote by  $k^+$  the maximal real subfield of  $k$  and by  $k_\infty^+$  the cyclotomic  $\mathbb{Z}_p$ -extension of  $k$ . If all of the Iwasawa invariants  $\lambda$ ,  $\mu$ , and  $\nu$  of  $k_\infty^+/k^+$  are zero (that is,  $\text{Gal}(L(k_\infty^+)/k_\infty^+)$  is trivial), then GGC holds for  $k$  and  $p$ .*

Here we shall give an example. Let  $k$  be the fifth cyclotomic field, and  $p$  a prime number which splits completely in  $k$  (that is,  $p \equiv 1 \pmod{5}$ ). Note that the class number of  $k$  is 1, and the maximal real subfield of  $k$  is  $\mathbb{Q}(\sqrt{5})$ . Assume that all of the Iwasawa invariants  $\lambda$ ,  $\mu$ , and  $\nu$  of the cyclotomic  $\mathbb{Z}_p$ -extension of  $\mathbb{Q}(\sqrt{5})$  are zero. Then GGC for  $k$  and  $p$  holds by Theorem 1.1. This assumption is satisfied for many primes  $p$ . (See, e.g., Sumida-Takahashi [23]. Many real quadratic fields and primes satisfying  $\lambda = \mu = \nu = 0$  are also given there.)

The outline of the proof is as follows. First, we take a certain  $\mathbb{Z}_p$ -extension  $K^{(1)}/k$  and a certain  $\mathbb{Z}_p^2$ -extension  $K^{(2)}/k$ . We show that  $\text{Gal}(L(K^{(1)})/K^{(1)})$  is trivial (Corollary 3.3). From this, we can see that  $\text{Gal}(L(K^{(2)})/K^{(2)})$  is pseudo-null as a  $\mathbb{Z}_p[[\text{Gal}(K^{(2)}/k)]]$ -module (Proposition 4.1). Finally, we will prove Theorem 1.1 by using these results. Our method can be considered as a generalization of the proof of Minardi's result which is mentioned above (see Sections 4 and 5).

## 2. Preliminaries

Let  $p$  be an odd prime number and  $F$  an algebraic number field. We denote by  $E(F)$  the group of units of  $F$ , and by  $A(F)$  the Sylow  $p$ -subgroup of the ideal class group of  $F$ . For a prime  $\mathfrak{p}$  of  $F$ , let  $U_{\mathfrak{p}}$  (resp.  $U_{\mathfrak{p}}^1$ ) be the group of units (resp. principal units) in the completion of  $F$  at  $\mathfrak{p}$ . Let  $F'$  be an algebraic extension of the field  $\mathbb{Q}$  of rational numbers. We write  $L(F')$  for the maximal unramified abelian pro- $p$  extension of  $F'$ , and  $\mathcal{M}_{\mathfrak{p}}(F')$  for the maximal pro- $p$  abelian extension of  $F'$  which is unramified outside the primes lying above  $\mathfrak{p}$  (where  $F$  is a finite extension of  $\mathbb{Q}$  contained in  $F'$ , and  $\mathfrak{p}$  is a prime of  $F$ ). If  $F'/F$  is a  $\mathbb{Z}_p$ -extension, we denote by  $\lambda(F'/F)$ ,  $\mu(F'/F)$ , and  $\nu(F'/F)$  the Iwasawa  $\lambda$ -invariant,  $\mu$ -invariant, and  $\nu$ -invariant of  $F'/F$ , respectively. Let  $T$  be a non-empty finite set of primes of  $F$  lying above  $p$  (it is not necessary that  $T$  contains all primes lying above  $p$ ). We put  $U_T = \prod_{\mathfrak{p} \in T} U_{\mathfrak{p}}$  and  $U_T^1 = \prod_{\mathfrak{p} \in T} U_{\mathfrak{p}}^1$ . We define the map  $\varphi: E(F) \rightarrow U_T$  by embedding diagonally. We denote by  $\mathcal{E}(F)_T$  the closure of  $\varphi(E(F)) \cap U_T^1$  in  $U_T^1$ . For the case that  $T = \{\mathfrak{p}\}$ , we will often write  $\mathcal{E}(F)_{\mathfrak{p}}$  instead of  $\mathcal{E}(F)_T$ .

The following result will be used frequently in our proof of the main theorem.

**Proposition 2.1.** (See Maire [14].) *Let  $k$  be an imaginary abelian quartic field and  $p$  an odd prime number which splits completely in  $k$ . We take a prime  $\mathfrak{P}$  of  $k$  lying above  $p$ . Let  $K$  be a finite abelian extension over  $k$ . Then  $\mathcal{M}_{\mathfrak{P}}(K)/K$  is a finite extension.*

**Proof.** This is a special case of [14, Theorem 25].  $\square$

**Lemma 2.2.** *Let  $k$  be an imaginary abelian quartic field and  $p$  an odd prime number which splits completely in  $k$ . For distinct primes  $\mathfrak{P}$  and  $\mathfrak{P}'$  of  $k$  lying above  $p$ , there exists a unique  $\mathbb{Z}_p$ -extension  $K/k$  which is unramified outside  $\{\mathfrak{P}, \mathfrak{P}'\}$ . Moreover, every prime of  $k$  lying above  $p$  is finitely decomposed in  $K$ .*

**Proof.** We put  $T = \{\mathfrak{P}, \mathfrak{P}'\}$ . Let  $\mathcal{M}_T(k)$  be the maximal abelian pro- $p$  extension over  $k$  which is unramified outside the primes contained in  $T$ . Then by class field theory, we have the following exact sequence (see [9, Proposition 3.1], [14, Theorem 5]):

$$0 \rightarrow U_T^1/\mathcal{E}(k)_T \rightarrow \text{Gal}(\mathcal{M}_T(k)/k) \rightarrow A(k) \rightarrow 0.$$

In this case, the  $\mathbb{Z}_p$ -rank of  $U_T^1$  is 2. Since the  $\mathbb{Z}$ -rank of  $E(k)$  is 1, the  $\mathbb{Z}_p$ -rank of  $\mathcal{E}(k)_T$  is also 1. We see that the  $\mathbb{Z}_p$ -rank of  $\text{Gal}(\mathcal{M}_T(k)/k)$  is 1 because  $A(k)$  is finite. The existence and the uniqueness have been proved.

We shall show the latter part. We note that both of  $\mathfrak{P}$  and  $\mathfrak{P}'$  must be ramified infinitely by Proposition 2.1. Hence both of  $\mathfrak{P}$  and  $\mathfrak{P}'$  are finitely decomposed. Next, we take an arbitrary prime  $\mathfrak{P}''$  of  $k$  which is lying above  $p$  but different from  $\mathfrak{P}$  and  $\mathfrak{P}'$ . Let  $\mathcal{M}'$  be the maximal abelian pro- $p$  extension over  $k$  which is unramified outside the primes contained in  $T$  and  $\mathfrak{P}''$  splits completely. It is sufficient to show that the  $\mathbb{Z}_p$ -rank of  $\text{Gal}(\mathcal{M}'/k)$  is zero. Let  $E(k)''$  be the group of  $\mathfrak{P}''$ -units in  $k$  and  $\varphi(E(k)'')$  the diagonal image of  $E(k)''$  in  $U_T$ . We denote by  $\mathcal{E}(k)_T''$  the closure of  $\varphi(E(k)'') \cap U_T^1$  in  $U_T^1$ . By class field theory, the  $\mathbb{Z}_p$ -rank of  $\text{Gal}(\mathcal{M}'/k)$  is equal to the  $\mathbb{Z}_p$ -rank of  $U_T^1/\mathcal{E}(k)_T''$  (see [14, Theorem 5]). Hence, we shall show that the  $\mathbb{Z}_p$ -rank of  $\mathcal{E}(k)_T''$  is exactly 2. Assume that the  $\mathbb{Z}_p$ -rank of  $\mathcal{E}(k)_T''$  is 1. Then there exists an element  $\pi$  of  $E(k)''$  which satisfies the following properties:

- $(\mathfrak{P}'')^h = (\pi)$  with a positive integer  $h$ , and
- for every positive integer  $n$ ,

$$\pi \equiv \varepsilon_n \pmod{\mathfrak{P}^n}, \quad \pi \equiv \varepsilon_n \pmod{\mathfrak{P}'^n}$$

with a unit  $\varepsilon_n$  of  $k$ .

It is known that the index  $(E(k) : W(k)E(k^+))$  is 1 or 2 (see, e.g. [24, Theorem 4.12]), where  $W(k)$  is the group of roots of unity in  $k$ . Hence we see that  $\varepsilon_n^{4w} \in k^+$  and  $N_{k^+/\mathbb{Q}}(\varepsilon_n^{4w}) = 1$  for all  $n$ , where  $w$  is the order of  $W(k)$ . We take an element  $\sigma$  of  $\text{Gal}(k/\mathbb{Q})$  which satisfies  $(\mathfrak{P}')^\sigma = \mathfrak{P}$ . If  $\sigma$  is contained in  $\text{Gal}(k/k^+)$ , then  $\pi^{4w} \equiv (\pi^{4w})^\sigma \pmod{\mathfrak{P}^n}$  for all  $n$  because  $\varepsilon_n^{4w} = (\varepsilon_n^{4w})^\sigma$ . Hence  $\pi^{4w} = (\pi^{4w})^\sigma$ . It is a contradiction because  $(\pi^\sigma) = ((\mathfrak{P}'')^\sigma)^h$  and  $(\mathfrak{P}'')^\sigma \neq \mathfrak{P}''$ . Otherwise, we can see that  $\pi^{4w}(\pi^{4w})^\sigma \equiv 1 \pmod{\mathfrak{P}^n}$  for all  $n$  because  $\varepsilon_n^{4w}(\varepsilon_n^{4w})^\sigma = 1$ . Hence  $\pi^{4w}(\pi^{4w})^\sigma = 1$ . It is also a contradiction since  $\pi$  generates a power of a prime lying above  $p$ . This shows that the  $\mathbb{Z}_p$ -rank of  $\mathcal{E}(k)_T$  is greater than 1, and then  $\text{Gal}(\mathcal{M}'/k)$  is a finite group. Hence  $\mathfrak{P}''$  cannot be infinitely decomposed. We have shown the lemma completely.  $\square$

From now on, we fix the following notation:

- $k$ : an imaginary abelian quartic number field;
- $k^+$ : the maximal real subfield of  $k$ ;

$p$ : an odd prime number which splits completely in  $k$ ;  
 $p_1, p_2$ : distinct primes of  $k^+$  lying above  $p$ ;  
 $\mathfrak{p}_i, \mathfrak{p}'_i$ : distinct primes of  $k$  lying above  $p_i$  ( $i = 1, 2$ ).

By Lemma 2.2, there exists a unique  $\mathbb{Z}_p$ -extension  $K^{(1)}/k$  which is unramified outside  $\{\mathfrak{p}_1, \mathfrak{p}_2\}$ . We can also show that there exists a unique  $\mathbb{Z}_p^2$ -extension  $K^{(2)}/k$  which is unramified outside  $\{\mathfrak{p}_1, \mathfrak{p}_2, \mathfrak{p}'_1\}$  by using the same method given in the proof of Lemma 2.2. Let  $\tilde{k}$  be the composite of all  $\mathbb{Z}_p$ -extensions of  $k$ . It is known that  $\tilde{k}/k$  is a  $\mathbb{Z}_p^3$ -extension (see, e.g., [24, Theorem 13.4]). Clearly, we can see that  $K^{(1)} \subset K^{(2)} \subset \tilde{k}$ .

The following result is also important for the proof of our main theorem. This is a special case of [20, p. 12, Lemme 4].

**Proposition 2.3.** (See Perrin-Riou [20].) *Let  $d$  be an integer which satisfies  $d \geq 2$ ,  $K/k$  a  $\mathbb{Z}_p^d$ -extension, and  $K'/k$  a  $\mathbb{Z}_p^{d-1}$ -extension which satisfies  $K' \subset K$ . Let  $M$  be a finitely generated torsion  $\mathbb{Z}_p[[\text{Gal}(K/k)]]$ -module and  $M_{\text{Gal}(K/K')}$  the  $\text{Gal}(K/K')$ -coinvariant quotient module of  $M$ . If  $M_{\text{Gal}(K/K')}$  is pseudo-null as a module over  $\mathbb{Z}_p[[\text{Gal}(K'/k)]]$ , then  $M$  is pseudo-null as a module over  $\mathbb{Z}_p[[\text{Gal}(K/k)]]$ .*

### 3. The $\mathbb{Z}_p$ -extension $K^{(1)}/k$

At first, we shall state the following known result.

**Theorem 3.1.** (See Fukuda and Komatsu [3], Hachimori [9].) *Let  $F$  be a real quadratic field,  $p$  an odd prime number which splits in  $F$ , and  $F_\infty$  the cyclotomic  $\mathbb{Z}_p$ -extension of  $F$ . We denote by  $\varepsilon_F$  the fundamental unit of  $F$  and we regard  $\varepsilon_F$  as an element of  $\mathbb{Z}_p$  (with a fixed embedding of  $k$  in the field  $\mathbb{Q}_p$  of  $p$ -adic numbers). Then,  $\lambda(F_\infty/F) = \mu(F_\infty/F) = \nu(F_\infty/F) = 0$  if and only if  $p$  does not divide the class number of  $F$  and  $\varepsilon_F^{p-1} \not\equiv 1 \pmod{p^2\mathbb{Z}_p}$ .*

We shall show a non-cyclotomic analog of a part of the above theorem.

**Proposition 3.2.** *If the class number of  $k$  is not divisible by  $p$ ,  $U_{\mathfrak{p}_1}^1 = \mathcal{E}(k)_{\mathfrak{p}_1}$ , and  $U_{\mathfrak{p}_2}^1 = \mathcal{E}(k)_{\mathfrak{p}_2}$ , then  $\lambda(K^{(1)}/k) = \mu(K^{(1)}/k) = \nu(K^{(1)}/k) = 0$ .*

**Proof.** We have the following exact sequence:

$$0 \rightarrow U_{\mathfrak{p}_1}^1/\mathcal{E}(k)_{\mathfrak{p}_1} \rightarrow \text{Gal}(\mathcal{M}_{\mathfrak{p}_1}(k)/k) \rightarrow A(k) \rightarrow 0$$

by class field theory (recall also the first part of the proof of Lemma 2.2). From the assumptions, we see that  $\text{Gal}(\mathcal{M}_{\mathfrak{p}_1}(k)/k)$  is trivial. Hence there is no abelian  $p$ -extension which is unramified outside  $\mathfrak{p}_1$ . This implies that  $\mathfrak{p}_2$  is totally ramified in  $K^{(1)}/k$ . Similarly, we can see that  $\mathfrak{p}_1$  is also totally ramified in  $K^{(1)}/k$ .

The rest of the proof can be shown by using the idea given in [9] (see also Chapter 13 of [24]). It is sufficient to show that  $L(K^{(1)})$  coincides with  $K^{(1)}$ . We note that  $\text{Gal}(\mathcal{M}_{\mathfrak{p}_1}(K^{(1)})/K^{(1)})$  is a compact  $\mathbb{Z}_p[[\text{Gal}(K^{(1)}/k)]]$ -module. Let  $I_{\mathfrak{p}_2}$  be the inertia group of  $\text{Gal}(\mathcal{M}_{\mathfrak{p}_1}(K^{(1)})/k)$  for some prime lying above  $\mathfrak{p}_2$ . We note that  $I_{\mathfrak{p}_2} \cap \text{Gal}(\mathcal{M}_{\mathfrak{p}_1}(K^{(1)})/K^{(1)})$  is trivial because  $\mathcal{M}_{\mathfrak{p}_1}(K^{(1)})/K^{(1)}$  is unramified outside the primes lying above  $\mathfrak{p}_1$ . Since  $\mathfrak{p}_2$  is totally ramified in  $K^{(1)}/k$ , we see that  $K^{(1)} \cap \mathcal{M}_{\mathfrak{p}_1}(k) = k$ . From these facts, we can see that the  $\text{Gal}(K^{(1)}/k)$ -coinvariant quotient of  $\text{Gal}(\mathcal{M}_{\mathfrak{p}_1}(K^{(1)})/K^{(1)})$  is isomorphic to  $\text{Gal}(\mathcal{M}_{\mathfrak{p}_1}(k)/k)$ . Since  $\text{Gal}(\mathcal{M}_{\mathfrak{p}_1}(k)/k)$  is trivial,  $\text{Gal}(\mathcal{M}_{\mathfrak{p}_1}(K^{(1)})/K^{(1)})$  is also trivial by Nakayama's lemma. We have shown the proposition because  $L(K^{(1)})$  is contained in  $\mathcal{M}_{\mathfrak{p}_1}(K^{(1)})$ .  $\square$

**Remark.** For the case that  $p = 2$  and  $k$  is biquadratic, a similar result is given in [12]. Goto [4] independently proved a similar result for more general situations.

Let  $k_{\infty}^{+}$  be the cyclotomic  $\mathbb{Z}_p$ -extension of the real quadratic field  $k^{+}$ . As a consequence of the above results, we have the following:

**Corollary 3.3.** *We assume that the class number of  $k$  is not divisible by  $p$  and  $\lambda(k_{\infty}^{+}/k^{+}) = \mu(k_{\infty}^{+}/k^{+}) = \nu(k_{\infty}^{+}/k^{+}) = 0$ . Then  $\lambda(K^{(1)}/k) = \mu(K^{(1)}/k) = \nu(K^{(1)}/k) = 0$ . (That is,  $\text{Gal}(L(K^{(1)})/K^{(1)})$  is trivial.)*

**Proof.** Since  $p$  splits completely in  $k$ , we see that  $U_{\mathfrak{P}_1}^1$  is isomorphic to  $U_{\mathfrak{p}_1}^1$ . By the assumption and Theorem 3.1, we obtain that  $U_{\mathfrak{p}_1}^1 = \mathcal{E}(k^{+})_{\mathfrak{p}_1}$ . Hence we see that  $U_{\mathfrak{P}_1}^1 = \mathcal{E}(k)_{\mathfrak{P}_1}$ . By using the same method, we also see that  $U_{\mathfrak{P}_2}^1 = \mathcal{E}(k)_{\mathfrak{P}_2}$ . The assertion follows from Proposition 3.2.  $\square$

#### 4. The $\mathbb{Z}_p^2$ -extension $K^{(2)}/k$

Recall that  $K^{(2)}$  is the unique  $\mathbb{Z}_p^2$ -extension of  $k$  which is unramified outside  $\{\mathfrak{P}_1, \mathfrak{P}_2, \mathfrak{P}'_1\}$ . We put  $X_1 = \text{Gal}(L(K^{(1)})/K^{(1)})$ ,  $X_2 = \text{Gal}(L(K^{(2)})/K^{(2)})$ , and  $G = \text{Gal}(K^{(2)}/K^{(1)}) \cong \mathbb{Z}_p$ .

**Proposition 4.1.** *Assume that neither  $\mathfrak{P}_1$  nor  $\mathfrak{P}_2$  splits in  $K^{(1)}/k$ . If the order of  $X_1$  is finite, then  $X_2$  is pseudo-null  $\mathbb{Z}_p[\text{Gal}(K^{(2)}/k)]$ -module.*

**Proof.** Our proof is quite similar to the proof of Minardi's result [16,17] (for the proof, it is convenient to see [19, §4]). However, we shall give a detailed proof because there are some points which must be confirmed carefully.

We denote by  $(X_2)_G$  the  $G$ -coinvariant quotient module of  $X_2$ . By Proposition 2.3, it is sufficient to show that the order of  $(X_2)_G$  is finite. Let  $L_1$  be the maximal subfield of  $L(K^{(2)})$  which is abelian over  $K^{(1)}$ . Then  $\text{Gal}(L_1/K^{(2)})$  is isomorphic to  $(X_2)_G$ .

By Lemma 2.2, the prime  $\mathfrak{P}'_1$  of  $k$  is finitely decomposed in  $K^{(1)}$ . Then we take a positive integer  $n_0$  such that the decomposition field of  $K^{(1)}/k$  for  $\mathfrak{P}'_1$  coincides with  $K_{n_0}^{(1)}$  (where  $K_{n_0}^{(1)}$  is the  $n_0$ th layer of  $K^{(1)}/k$ ).

We put  $\mathcal{I}_1 = \sum_P I_P$ , where  $P$  runs all primes of  $K^{(1)}$  which ramify in  $L_1/K^{(1)}$ , and  $I_P$  denotes the inertia group of  $\text{Gal}(L_1/K^{(1)})$  for  $P$ . We have the following exact sequence of  $\mathbb{Z}_p[\text{Gal}(K^{(1)}/k)]$ -modules:

$$0 \rightarrow \mathcal{I}_1 \rightarrow \text{Gal}(L_1/K^{(1)}) \rightarrow \text{Gal}(L(K^{(1)})/K^{(1)}) \rightarrow 0.$$

We note that all primes of  $K^{(1)}$  which ramify in  $L_1/K^{(1)}$  are the primes lying above  $\mathfrak{P}'_1$ . We also note that  $\mathfrak{P}'_1$  is finitely decomposed in  $K^{(1)}$  by Lemma 2.2. Hence  $\mathcal{I}_1$  is finitely generated over  $\mathbb{Z}_p$  because each  $I_P$  is isomorphic to the additive group of  $\mathbb{Z}_p$ . This implies that  $\mathcal{I}_1$  is a finitely generated torsion  $\mathbb{Z}_p[\text{Gal}(K^{(1)}/k)]$ -module. By the assumption that  $X_1 = \text{Gal}(L(K^{(1)})/K^{(1)})$  is finite, we also see that  $\text{Gal}(L_1/K^{(1)})$  is a finitely generated torsion  $\mathbb{Z}_p[\text{Gal}(K^{(1)}/k)]$ -module, and  $\mathcal{I}_1$  is pseudo-isomorphic to  $\text{Gal}(L_1/K^{(1)})$ . Let  $L'_1$  be the subfield of  $L_1$  corresponding to the maximal pseudo-null (that is, finite) submodule of  $\text{Gal}(L_1/K^{(1)})$ .

We claim that  $L'_1$  is an abelian extension over  $K_{n_0}^{(1)}$ . Take an arbitrary prime  $P_1$  of  $K^{(1)}$  lying above  $\mathfrak{P}'_1$ , and let  $I_{P_1}$  be the inertia group of  $\text{Gal}(L'_1/K^{(1)})$  for  $P_1$ . Then  $\text{Gal}(K^{(1)}/K_{n_0}^{(1)})$  acts on  $I_{P_1}$ . Since  $L_1/K^{(2)}$  is an unramified extension, the restriction map  $I_{P_1} \rightarrow \text{Gal}(K^{(2)}/K^{(1)})$  is injective. We see that  $\text{Gal}(K^{(1)}/K_{n_0}^{(1)})$  acts trivially on  $I_{P_1}$  for each prime  $P_1$  lying above  $\mathfrak{P}'_1$ . Then,  $\text{Gal}(K^{(1)}/K_{n_0}^{(1)})$  acts trivially on  $\mathcal{I}_1$ . Since pseudo-isomorphism satisfies the symmetric law for torsion  $\mathbb{Z}_p[\text{Gal}(K^{(1)}/k)]$ -modules (see [18,24]) and  $\text{Gal}(L'_1/K^{(1)})$  has no nontrivial pseudo-null submodule, there is an injection

(as  $\mathbb{Z}_p[\text{Gal}(K^{(1)}/k)]$ -modules) from  $\text{Gal}(L'_1/K^{(1)})$  to  $\mathcal{I}_1$ . Hence  $\text{Gal}(K^{(1)}/K_{n_0}^{(1)})$  also acts trivially on  $\text{Gal}(L'_1/K^{(1)})$ . The claim follows.

By the assumption, neither  $\mathfrak{P}_1$  nor  $\mathfrak{P}_2$  splits in  $K^{(1)}/k$ . We denote by  $I_{\mathfrak{P}_1}$  (resp.  $I_{\mathfrak{P}_2}$ ) the inertia group of  $L'_1/K_{n_0}^{(1)}$  for the prime of  $K_{n_0}^{(1)}$  lying above  $\mathfrak{P}_1$  (resp.  $\mathfrak{P}_2$ ). Note that the  $\mathbb{Z}_p$ -rank of  $I_{\mathfrak{P}_1}$  (resp.  $I_{\mathfrak{P}_2}$ ) is 1 because the prime of  $K^{(1)}$  lying above  $\mathfrak{P}_1$  (resp.  $\mathfrak{P}_2$ ) is unramified in  $L'_1$ . Let  $M_1$  be the fixed field of  $L'_1$  by  $\sum_{i=1,2} I_{\mathfrak{P}_i}$ . Then we have the following exact sequence:

$$0 \rightarrow \sum_{i=1,2} I_{\mathfrak{P}_i} \rightarrow \text{Gal}(L'_1/K_{n_0}^{(1)}) \rightarrow \text{Gal}(M_1/K_{n_0}^{(1)}) \rightarrow 0.$$

Since  $M_1$  is an abelian extension of  $K_{n_0}^{(1)}$  which is unramified outside the primes lying above  $\mathfrak{P}'_1$ , we see that the order of  $\text{Gal}(M_1/K_{n_0}^{(1)})$  is finite by Proposition 2.1. Hence, we can conclude that the  $\mathbb{Z}_p$ -rank of  $\text{Gal}(L'_1/K_{n_0}^{(1)})$  is exactly 2. This implies that  $L'_1$  is a finite extension over  $K^{(2)}$  because the  $\mathbb{Z}_p$ -rank of  $\text{Gal}(K^{(2)}/K_{n_0}^{(1)})$  is 2. Since  $L_1/L'_1$  is a finite extension,  $L'_1$  is also a finite extension over  $K^{(2)}$ . Consequently, the order of  $(X_2)_G$  is finite, and then the proposition follows.  $\square$

## 5. Proof of Theorem 1.1

Let the notation be as in the previous sections. Recall that  $k$  is an imaginary abelian quartic field and  $p$  is an odd prime which splits completely in  $k$ . We assume that the class number of  $k$  is not divisible by  $p$ , and  $\lambda(k_{\infty}^+/k^+) = \mu(k_{\infty}^+/k^+) = \nu(k_{\infty}^+/k^+) = 0$ . By the results given in Section 3, we see that  $\mathfrak{P}_1$  and  $\mathfrak{P}_2$  are totally ramified in  $K^{(1)}/k$  and  $X_1$  is trivial. Hence  $X_2$  is pseudo-null as a  $\mathbb{Z}_p[\text{Gal}(K^{(2)}/k)]$ -module by Proposition 4.1.

The outline of the following argument is the same as the proof of Proposition 4.1. We put  $\tilde{X} = \text{Gal}(L(\tilde{k})/\tilde{k})$  and  $G_2 = \text{Gal}(\tilde{k}/K^{(2)})$ . We will show that  $G_2$ -coinvariant quotient  $(\tilde{X})_{G_2}$  is pseudo-null. Then by Proposition 2.3, the theorem follows.

We shall recall the argument given in [5]. Let  $L_2$  be the maximal subfield of  $L(\tilde{k})$  which is abelian over  $K^{(2)}$ . Then  $\text{Gal}(L_2/\tilde{k})$  is isomorphic to  $(\tilde{X})_{G_2}$ . For a prime  $P_2$  of  $K^{(2)}$  lying above  $\mathfrak{P}'_2$ , let  $I_{P_2}$  be the inertia group of  $L_2/K^{(2)}$  for  $P_2$ . ( $I_{P_2}$  is isomorphic to the additive group of  $\mathbb{Z}_p$ .) Moreover, we put  $\mathcal{I}_2 = \sum_{P_2 \in \mathfrak{P}'_2} I_{P_2}$ . (Note that there are infinitely many primes in  $K^{(2)}$  lying above  $\mathfrak{P}'_2$ .) Since  $L_2/K^{(2)}$  is unramified outside the primes lying above  $\mathfrak{P}'_2$ , we have the following exact sequence of  $\mathbb{Z}_p[\text{Gal}(K^{(2)}/k)]$ -modules:

$$0 \rightarrow \mathcal{I}_2 \rightarrow \text{Gal}(L_2/K^{(2)}) \rightarrow \text{Gal}(L(K^{(2)})/K^{(2)}) \rightarrow 0. \quad (1)$$

Note that  $\mathcal{I}_2$  is generated by the inertia groups for the primes lying above  $\mathfrak{P}'_2$ , and these inertia groups are conjugate under the action of  $\text{Gal}(K^{(2)}/k)$ . Then  $\mathcal{I}_2$  is a finitely generated  $\mathbb{Z}_p[\text{Gal}(K^{(2)}/k)]$ -module. We see that  $\text{Gal}(L_2/K^{(2)})$  is also a finitely generated  $\mathbb{Z}_p[\text{Gal}(K^{(2)}/k)]$ -module. (These facts are essentially given in [5]. See the proof of [5, Theorem 1].)

Let  $D_2$  be the decomposition group of  $K^{(2)}/k$  for  $\mathfrak{P}'_2$ . We can see that  $D_2$  is nontrivial and isomorphic to the additive group of  $\mathbb{Z}_p$  by using Lemma 2.2. Take a topological generator  $\gamma$  of  $D_2$ . Let  $N$  be the fixed field of  $K^{(2)}$  by  $D_2$ , then we can see that there exists a positive integer  $n_1$  such that  $N/K_{n_1}^{(1)}$  is a  $\mathbb{Z}_p$ -extension (that is,  $K_{n_1}^{(1)}$  is the decomposition field of  $K^{(1)}/k$  for  $\mathfrak{P}'_2$ ).

We note that the  $\mathbb{Z}_p[\text{Gal}(K^{(2)}/k)]$ -modules which appeared in (1) can be seen as modules over  $\mathbb{Z}_p[\text{Gal}(K^{(2)}/K_{n_1}^{(1)})]$ . Here, we will show that  $(\tilde{X})_{G_2}$  is pseudo-null as a  $\mathbb{Z}_p[\text{Gal}(K^{(2)}/K_{n_1}^{(1)})]$ -module. Fix a topological generator  $\sigma$  of  $\text{Gal}(K^{(2)}/K^{(1)})$ . We can show that  $\text{Gal}(K^{(2)}/K_{n_1}^{(1)}) \cong \mathbb{Z}_p^2$  is topologically generated by  $\sigma$  and  $\gamma$ . Take an isomorphism from  $\mathbb{Z}_p[\text{Gal}(K^{(2)}/K_{n_1}^{(1)})]$  to  $\Lambda_2 = \mathbb{Z}_p[[T_1, T_2]]$  which satisfies  $\sigma \mapsto 1 + T_1$  and  $\gamma \mapsto 1 + T_2$ . We shall identify  $\mathbb{Z}_p[\text{Gal}(K^{(2)}/K_{n_1}^{(1)})]$  and  $\Lambda_2$  via this isomorphism. (For a detailed theory of  $\Lambda_2$ -modules, see, e.g., [7,18].)

Since the restriction map  $I_{P_2} \rightarrow \text{Gal}(\tilde{k}/K^{(2)})$  is injective, we see  $\text{Gal}(K^{(2)}/N)$  acts trivially on  $I_{P_2}$  for each prime  $P_2$  lying above  $\mathfrak{P}'_2$ . This implies that  $\gamma - 1$  annihilates  $\mathcal{I}_2$ . Then  $\mathcal{I}_2$  is a finitely generated torsion  $\Lambda_2$ -module. We remark that  $X_2$  is a finitely generated torsion  $\Lambda_2$ -module. Moreover,  $X_2$  is also pseudo-null as a  $\Lambda_2$ -module. This follows from, for example, [13, Lemma 2.3] and an equivalent condition of pseudo-nullity (see the second paragraph of Section 1). (The same fact is also mentioned in [16].) Then by the exact sequence (1), we see that  $\text{Gal}(L_2/K^{(2)})$  is a finitely generated torsion  $\Lambda_2$ -module and  $\mathcal{I}_2$  is pseudo-isomorphic to  $\text{Gal}(L_2/K^{(2)})$ . (Note that both of  $\mathcal{I}_2$  and  $\text{Gal}(L_2/K^{(2)})$  are also finitely generated torsion  $\mathbb{Z}_p[[\text{Gal}(K^{(2)}/k)]]$ -modules.)

Let  $L'_2$  be the subfield of  $L_2$  corresponding to the maximal pseudo-null submodule (as a  $\Lambda_2$ -module) of  $\text{Gal}(L_2/K^{(2)})$ . Since  $\mathcal{I}_2$  is pseudo-isomorphic to  $\text{Gal}(L_2/K^{(2)})$ , we can see that  $\gamma$  acts trivially on  $\text{Gal}(L'_2/K^{(2)})$  by using the same argument given in the proof of Proposition 4.1 (see the fifth paragraph of the proof of Proposition 4.1). Hence we see that  $L'_2$  is an abelian extension over  $N$ . Note that  $\text{Gal}(N/K_{n_1}^{(1)})$  acts on  $\text{Gal}(L'_2/K^{(2)})$  because  $\text{Gal}(L'_2/K^{(2)})$  is a module over  $\mathbb{Z}_p[[\text{Gal}(K^{(2)}/k)]]$ .

**Lemma 5.1.**  $L'_2/N$  is unramified outside the primes lying above  $\mathfrak{P}'_2$ .

**Proof.** Let  $N'$  be the unique  $\mathbb{Z}_p$ -extension of  $k$  contained in  $N$ . It is sufficient to show that all of  $\mathfrak{P}_1$ ,  $\mathfrak{P}_2$ , and  $\mathfrak{P}'_1$  are ramified in  $N'/k$  because all of the inertia groups of  $K^{(2)}/k$  for  $\mathfrak{P}_1$ ,  $\mathfrak{P}_2$ , and  $\mathfrak{P}'_1$  are isomorphic to  $\mathbb{Z}_p$ , and  $L'_2/K^{(2)}$  is unramified outside the primes lying above  $\mathfrak{P}'_2$ .

By Proposition 2.1, there is no  $\mathbb{Z}_p$ -extension of  $k$  in which only one prime of  $k$  lying above  $p$  ramifies. Then we assume that there are only two primes in  $k$  which are ramified in  $N'/k$ . However, it is already seen in Lemma 2.2 that every prime of  $k$  lying above  $p$  is finitely decomposed in  $N'$ . It is a contradiction because  $\mathfrak{P}'_2$  is decomposed completely in  $N'$ .  $\square$

**Lemma 5.2.**  $\text{Gal}(L'_2/K^{(2)})$  is a finitely generated torsion  $\mathbb{Z}_p[[\text{Gal}(N/K_{n_1}^{(1)})]]$ -module.

**Proof.** By Lemma 5.1, we see that  $L'_2$  is contained in  $\mathcal{M}_{\mathfrak{P}'_2}(N)$ . We shall show that  $\text{Gal}(\mathcal{M}_{\mathfrak{P}'_2}(N)/N)$  is a finitely generated torsion  $\mathbb{Z}_p[[\text{Gal}(N/K_{n_1}^{(1)})]]$ -module. Then the lemma follows from this claim.

Because  $N/K_{n_1}^{(1)}$  is a  $\mathbb{Z}_p$ -extension, we can identify  $\mathbb{Z}_p[[\text{Gal}(N/K_{n_1}^{(1)})]]$  as the one variable power series ring  $\mathbb{Z}_p[[T]]$ . For a positive integer  $m$ , let  $N_m$  be the  $m$ th layer of  $N/K_{n_1}^{(1)}$  and  $\mathcal{M}_{\mathfrak{P}'_2}(N_m)$  the maximal abelian pro- $p$  extension of  $N_m$  which is unramified outside the primes lying above  $\mathfrak{P}'_2$ . By using a well-known argument of Iwasawa theory (see, e.g., [9,24]), there exists a positive integer  $m_0$  such that for all  $m > m_0$ ,

$$\text{Gal}(\mathcal{M}_{\mathfrak{P}'_2}(N_m)/N_m) \cong \text{Gal}(\mathcal{M}_{\mathfrak{P}'_2}(N)/N)/v_{m,m_0}W,$$

where

$$W = \text{Gal}(\mathcal{M}_{\mathfrak{P}'_2}(N)/N\mathcal{M}_{\mathfrak{P}'_2}(N_{m_0}))$$

and

$$v_{m,m_0} = \frac{(1+T)^{p^m} - 1}{(1+T)^{p^{m_0}} - 1} \in \mathbb{Z}_p[[T]].$$

By Proposition 2.1, we can see that  $\text{Gal}(\mathcal{M}_{\mathfrak{P}'_2}(N_m)/N_m)$  is finite for all  $m$ . Using the argument given in Chapter 13 of [24] or [9, Lemma 6.1], we can show that  $\text{Gal}(\mathcal{M}_{\mathfrak{P}'_2}(N)/N)$  is a finitely generated torsion module over  $\mathbb{Z}_p[[\text{Gal}(N/K_{n_1}^{(1)})]]$ .  $\square$

We shall finish the proof of Theorem 1.1. Recall that  $\sigma$  is a topological generator of  $\text{Gal}(K^{(2)}/K^{(1)})$  and  $\gamma$  is a topological generator of  $\text{Gal}(K^{(2)}/N)$ . Recall also that we fixed an isomorphism

$\mathbb{Z}_p[\text{Gal}(K^{(2)}/K_{n_1}^{(1)})]$  to  $\Lambda_2$  such that  $\sigma \mapsto 1 + T_1$  and  $\gamma \mapsto 1 + T_2$ . Since  $L'_2/N$  is an abelian extension, we see that  $T_2$  annihilates  $\text{Gal}(L'_2/K^{(2)})$ . By Lemma 5.2 and the fact that  $\text{Gal}(N/K_{n_1}^{(1)}) \cong \text{Gal}(K^{(2)}/K^{(1)})$ , we can take a nonzero element  $f(T_1)$  of  $\mathbb{Z}_p[[T_1]]$  such that  $f(T_1)$  annihilates  $\text{Gal}(L'_2/K^{(2)})$ . Since  $f(T_1)$  and  $T_2$  are relatively prime,  $\text{Gal}(L'_2/K^{(2)})$  is annihilated by two relatively prime elements of  $\Lambda_2$ . Hence we showed that  $\text{Gal}(L'_2/K^{(2)})$  is a pseudo-null  $\Lambda_2$ -module.

By the definition of  $L'_2$  (see the paragraph before Lemma 5.1), we see that  $\text{Gal}(L_2/K^{(2)})$  is a pseudo-null  $\Lambda_2$ -module. Therefore, we conclude that  $(\tilde{X})_{G_2} \cong \text{Gal}(L_2/\tilde{k})$  is a pseudo-null  $\Lambda_2$ -module. By [13, Lemma 2.3] and an equivalent condition of pseudo-nullity (see the second paragraph of Section 1), we see that  $(\tilde{X})_{G_2}$  is also pseudo-null as a  $\mathbb{Z}_p[[\text{Gal}(K^{(2)}/k)]]$ -module. Our main theorem has been proved.

**Remark.** We can prove our main theorem under a weaker condition. That is, if neither  $\mathfrak{P}_1$  nor  $\mathfrak{P}_2$  splits in  $K^{(1)}/k$  and  $X_1$  is finite, then GGC holds for  $k$  and  $p$ . This follows from Proposition 4.1 and the argument given in Section 5. This result is used in [11].

## Acknowledgments

The author express his thanks to Professor Norio Adachi for his advice and encouragement, to Professor Keiichi Komatsu for giving helpful comments, to Professor Manabu Ozaki for giving valuable remarks about the theory of multiple  $\mathbb{Z}_p$ -extensions, and to Professor Hiroki Sumida-Takahashi for giving comments about the example stated after Theorem 1.1. The author also express his thanks to the referee for giving many useful comments to the manuscript. The main part of this work was done while the author was in Waseda University, and this work was supported by Waseda University Grand for Special Research Projects (2003A-081).

## References

- [1] A. Bandini, Greenberg's conjecture for  $\mathbb{Z}_p^d$ -extensions, *Acta Arith.* 108 (2003) 357–368.
- [2] A. Bandini, Greenberg's conjecture and capitulation in  $\mathbb{Z}_p^d$ -extensions, *J. Number Theory* 122 (2007) 121–134.
- [3] T. Fukuda, K. Komatsu, On  $\mathbb{Z}_p$ -extensions of real quadratic fields, *J. Math. Soc. Japan* 38 (1986) 95–102.
- [4] H. Goto, Iwasawa invariants on non-cyclotomic  $\mathbb{Z}_p$ -extensions of CM fields, *Proc. Japan Acad. Ser. A Math. Sci.* 82 (2006) 152–154.
- [5] R. Greenberg, The Iwasawa invariants of  $\Gamma$ -extensions of a fixed number field, *Amer. J. Math.* 95 (1973) 204–214.
- [6] R. Greenberg, On the Iwasawa invariants of totally real number fields, *Amer. J. Math.* 98 (1976) 263–284.
- [7] R. Greenberg, On the structure of certain Galois groups, *Invent. Math.* 47 (1978) 85–99.
- [8] R. Greenberg, Iwasawa theory—past and present, in: *Class Field Theory—Its Centenary and Prospect*, in: *Adv. Stud. Pure Math.*, vol. 30, Math. Soc. Japan, Tokyo, 2001, pp. 335–385.
- [9] Y. Hachimori, Iwasawa theory for extensions with restricted  $p$ -ramification, *Tokyo J. Math.* 26 (2003) 179–198.
- [10] T. Itoh, Pseudo-null Iwasawa modules for  $\mathbb{Z}_2^2$ -extensions, *Tokyo J. Math.* 30 (2007) 199–209.
- [11] T. Itoh, Some questions on the ideal class group of imaginary abelian fields, *Math. J. Okayama Univ.* 49 (2007) 185–196.
- [12] T. Itoh, S. Kiyofuji, Y. Mizusawa, On the Iwasawa invariants of certain non-cyclotomic  $\mathbb{Z}_2$ -extensions, *Manuscripta Math.* 118 (2005) 227–233.
- [13] U. Jannsen, Iwasawa modules up to isomorphism, in: *Algebraic Number Theory—In Honor of K. Iwasawa*, in: *Adv. Stud. Pure Math.*, vol. 17, Kinokuniya/Academic Press, Tokyo/Boston, MA, 1989, pp. 171–207.
- [14] C. Maire, On the  $\mathbb{Z}_l$ -rank of abelian extensions with restricted ramification, *J. Number Theory* 92 (2002) 376–404.
- [15] W.G. McCallum, R.T. Sharifi, A cup product in the Galois cohomology of number fields, *Duke Math. J.* 120 (2003) 269–310.
- [16] J. Minardi, Iwasawa modules for  $\mathbb{Z}_p^d$ -extensions of algebraic number fields, thesis, University of Washington, 1986.
- [17] J. Minardi, Iwasawa modules for  $\mathbb{Z}_p^d$ -extensions of number fields, in: *Number Theory*, in: *CMS Conf. Proc.*, vol. 7, Amer. Math. Soc., Providence, RI, 1987, pp. 237–242.
- [18] J. Neukirch, A. Schmidt, K. Wingberg, *Cohomology of Number Fields*, second ed., *Grundlehren Math. Wiss.*, vol. 323, Springer-Verlag, Berlin, Heidelberg, 2008.
- [19] M. Ozaki, Iwasawa invariants of  $\mathbb{Z}_p$ -extensions over an imaginary quadratic field, in: *Class Field Theory—Its Centenary and Prospect*, in: *Adv. Stud. Pure Math.*, vol. 30, Math. Soc. Japan, Tokyo, 2001, pp. 387–399.
- [20] B. Perrin-Riou, Arithmétique des courbes elliptiques et théorie d'Iwasawa, *Mém. Soc. Math. France (N.S.)* 17 (1984) 1–130.
- [21] R.T. Sharifi, Iwasawa theory and the Eisenstein ideal, *Duke Math. J.* 137 (2007) 63–101.
- [22] R.T. Sharifi, On Galois groups of unramified pro- $p$  extensions, *Math. Ann.* 342 (2008) 297–308.
- [23] H. Sumida-Takahashi, The Iwasawa invariants and the higher  $K$ -groups associated to real quadratic fields, *Experiment. Math.* 14 (2005) 307–316.
- [24] L.C. Washington, *Introduction to Cyclotomic Fields*, second ed., *Grad. Texts in Math.*, vol. 83, Springer-Verlag, New York, Berlin, Heidelberg, 1997.